## Childhood and Adolescent Immunization Schedule • New Mexico • 2013-2014

Age ▶ Vaccine ▼	Birth	1 mo	2 mo	3 mo	4 mo	5 mo	6 mo	7 mo	8 mo	9 mo	10 mo	11 mo	12 mo (1 yr)	18 mo (1½yrs)	24 mo (2 yrs)	48 mo (4 yrs)	60 mo (5 yrs)	11-12 Yrs	13-18 Yrs
DTaP (Diphtheria- Tetanus-Pertussis)			Dose 1		Dose 2		Dose 3 and 4 should be 6 months apart					Dose 4			Dose 5				
Haemophilus Influenza B (Hib)			Dose 1		Dose 2		Dose 3*						Dose 4						
Hepatitis A							Dose Dose 1 2								Doses 1 and 2 should be 6 months apart				
Hepatitis B	At Birth	1 1000					Dose 3 – Between 6 and 18 mor						onths						
Influenza							Yearly, ages 6 months to 18 years during the flu season												
Measles, Mumps & Rubella													Dose 1			Dose 2			
Pneumonia (PCV)			Dose 1		Dose 2		Dose Note: First 3 doses must be 2 months apart						Dose 4						
Polio			Dose 1		Dose 2		Dose 3									Dose 4			
Rotavirus			Dose 1		Dose 2		Dose 3 If Rotarix given at 2 and 4 mos, 3 <sup>rd</sup> dose not needed												
Varicella (Chicken Pox)													Dose 1			Dose 2			
Human Papilloma Virus (HPV)																		3 Doses**	
Meningitis (Meningococcol-MCV4)																		Dose 1	Dose 2***
Tdap (Tetanus- Diphtheria-Pertussis)							2										140	Booster	

<sup>\*</sup>Hib B: If PRP-OMP is administered at ages 2 and 4 months, a dose at age 6 months is not indicated. Hibrex should only be used for the booster dose in children aged 12 months – 4 years.

Presbyterian Health Plan recommends the Center for Disease Control and Prevention Immunization Schedule which is updated annually. If you have any questions you can access the CDC website at <a href="http://www.cdc.gov/vaccines/schedules/index.html">http://www.cdc.gov/vaccines/schedules/index.html</a>. You can also contact New Mexico Immunization Hotline toll free at 1-866-681-5872 or online at <a href="http://www.immunizenm.org/sched.shtml">http://www.immunizenm.org/sched.shtml</a>. Services are funded in part under contract with the State of New Mexico.



Shots can be given even when your child has a minor illness. If your child is not up-to-date, your child's Primary Care Practitioner can make a plan to make sure all shots are complete. Take your child's shot record with you to every healthcare visit.

Your child's Primary Care Practitioner may adjust this schedule based on what is best for your child.

Remember, your child is not protected against these diseases until all doses have been given.

Presbyterian exists to improve the health of the patients, members, and communities we serve.

For more information on immunizations, go to www.phs.org/shots



<sup>\*\*</sup>HPV: Dose 2 should be given 2 months after dose 1. Dose 3 should be given 6 months after dose 1.

<sup>\*\*\*</sup> MCV4: Dose 2 should be given 5 years after dose 1.